



Nestin

Data Sheet

Catalog Number:	MO15012	Host:	Mouse
Product Type:	Mouse Monoclonal IgG1 Protein G purified Cell Culture Supernatant. Clone: 196908	Species Reactivity:	Human, Primate
Immunogen Sequence:	human Nestin fragment (amino acid residues 618 - 1618) expressing NSO cells	Format:	Liquid 1mg/ml Solution in phosphate-buffered saline (PBS) with 5% Trehlose.
Applications:	Immunocytochemistry: 5-10 µg/mL Flow Cytometry-see: PLoS ONE 8(6): e68519. doi:10.1371/journal.pone.0068519. Dilutions listed as a recommendation. Optimal dilution should be determined by investigator.		
Storage:	Antibody can also be aliquotted and stored frozen at -20° C to -70° C in a manual defrost freezer for six months without detectable loss of activity. The antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Avoid repeated freeze-thaw cycles.		
References:	Turac G, Hindley CJ, Thomas R, Davis JA, Deleidi M, et al. (2013) Combined Flow Cytometric Analysis of Surface and Intracellular Antigens Reveals Surface Molecule Markers of Human Neurogenesis. PLoS ONE 8(6): e68519. doi:10.1371/journal.pone.0068519. Fabien G. Lafaille, Itai M. Pessach, Shen-Ying Zhang, Michael J. Ciancanelli, Melina Herman, Avinash Abhyankar, Shui-Wang Ying, Sotirios Keros, Peter A. Goldstein, Gustavo Mostoslavsky, Jose Ordovas-Montanes, Emmanuelle Jouanguy, Sabine Plancoulaine, Edmund Tu, Yechiel Elkabetz, Saleh Al-Muhsen, Marc Tardieu, Thorsten M. Schlaeger, George Q. Daley, Laurent Abel, Jean-Laurent Casanova, Lorenz Studer, Luigi D. Notarangelo. Impaired intrinsic immunity to HSV-1 in human iPSC-derived TLR3-deficient CNS cells. Nature (2012) doi:10.1038/nature11583. J. Simon Lunn, Crystal Pacut, Emily Stern, Stacey A. Sakowski, J. Matthew Velke, Sue O'Shea, Eva L. Feldman. Intrasplinal transplantation of neurogenin-expressing stem cells generates spinal cord neural progenitors. Neurobiology of Disease. Volume 46, Issue 1, April 2012, Pages 59–68.		


Application Notes

Immunocytochemistry

This antibody can be used with the appropriate secondary reagents at the concentration of 5 - 10 µg/mL in fixed cells. Cells were fixed with 4% paraformaldehyde and 0.15% picric acid in PBS at room temperature for 20 min., followed by blocking with PBS containing 10% normal donkey serum and 1% BSA at room temperature for 45 min. After blocking, cells were incubated with diluted primary antibody overnight at 4° C and followed by incubation for 1 hour with appropriate secondary antibody at room temperature. Between each step, cells were washed with PBS and 0.1% BSA.


FOR RESEARCH USE ONLY

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. V2-06/2013



Images: Immunostaining of [STEMEZ\(TM\) hNP1 Progenitors](#) before and after differentiation. The culture was highly homogenous with neural rosettes. (A) Neural rosettes (white arrows, bright field) from WA09 cells. A similar result also obtained with BG02 cell line. (B) Shown here, a neural rosette stained with *Nestin* (NES) (green) antibody. Propagated Neural progenitors showed expression of marker genes, *NES* (C) and *Musashi 1* (D) but not *SOX2* (E). Further differentiation produced neurons (*TUJ1*) (F), astrocytes (*GFAP*) (G) and oligodendrocytes (*myelin basic protein*) (H) (lower panel). DAPI (blue) was used for staining the nuclei (scale bar for (A) through (E) is 100 μ m and for the remaining figures 10 μ m). Differentiation (2007) DOI: [10.1111/j.1432-0436.2007.00256.x](#)...Dilutions: [NES](#) (1:100, Neuromics, Edina, MN), [MSI1](#) (1:100, Neuromics), [Tuj1](#) (1:500, Neuromics)...

Images: Accurate detection of intracellular antigens with optimized fixation-permeabilization conditions preserving surface antigens. Flow cytometric detection of TUJ1, MAP2 and nestin antigens in BJ fibroblasts and the neural SH-SY5Y cell line (A). TUJ1 and nestin are present in both cell lines, while the mature neuronal marker MAP2 was only detected in SH-SY5Y cells (arrows). Note stable fluorescent levels of the negative population, indicating low background staining using this protocol. Representative experiment of three independent repeats shown. (B) Corresponding validation by immunofluorescence analysis. (C) Quantitation of TUJ1, MAP2 and nestin intracellular antigen detection ($n=3$). Error bars indicate standard deviation. (D) Response of TUJ1 and MAP2 intracellular antigen expression to 6 DIV of 10 μ M retinoic acid (RA) treatment of SH-SY5Y cells. Note disappearance/reduction of subsets negative for these markers (upward shift, green arrows), as well as a shift toward CD184^{low} expression with differentiation (blue arrows). doi:10.1371/journal.pone.0068519.g003



FOR RESEARCH USE ONLY

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. V2-06/2013

Gentaur Molecular Products
Voortstraat 49
1910 Kampenhout, BELGIUM

Tel 0032 16 58 90 45 | Fax 0032 16 50 90 45

www.gentaur-worldwide.com

info@gentaur.com